



The 1-h post-load plasma glucose as a novel biomarker for diagnosing dysglycemia

Ram Jagannathan¹ · Martin Buysschaert² · José Luis Medina³ · Karin Katz⁴ · Sarah Musleh⁴ · Brenda Dorcely⁴ · Michael Bergman⁴

Received: 24 August 2017 / Accepted: 11 January 2018
© Springer-Verlag Italia S.r.l., part of Springer Nature 2018

Abstract

Identifying the earliest moment for intervention to avert progression to prediabetes and diabetes in high-risk individuals is a substantial challenge. As β -cell function is already compromised in prediabetes, attention should therefore be focused on identifying high-risk individuals earlier in the so-called pre-prediabetes stage. Biomarkers to monitor progression and identify the time point at which β -cell dysfunction occurs are therefore critically needed. Large-scale population studies have consistently shown that the 1-h plasma glucose (1-h PG) ≥ 155 mg/dl (8.6 mmol/l) during the oral glucose tolerance test detected incident type 2 diabetes and associated complications earlier than fasting plasma glucose or 2-h plasma glucose levels. An elevated 1-h PG level appears to be a better alternative to HbA1c [5.7–6.4% (37–47 mmol/mol)] or traditional glucose criteria for identifying high-risk individuals at a stage when β -cell function is substantially more intact than in prediabetes. Diagnosing high-risk individuals earlier proffers the opportunity for potentially reducing progression to diabetes, development of microvascular complications and mortality, thereby advancing benefit beyond that which has been demonstrated in global diabetes prevention programs.

Keywords Dysglycemia · Prediabetes · Diabetes · Oral glucose tolerance test

Introduction

The International Diabetes Federation (IDF) estimates that globally 425 million individuals, or 8.8% (1 in 11 adults) have diabetes (T2DM) with 629 million adults or 9.9% of the world's population expected to develop diabetes by 2045 [1]. In addition, 7.3% of the world's population or 352 million

have impaired glucose tolerance (IGT) and are considered at high risk for developing diabetes with an expectation that this will increase to 532 million, or 8.3%, in 2045 [1]. Forty percent of adults face a lifetime risk of diabetes representing a substantial increase from 20% in the late 1980s [2]. Delaying the diagnosis can result in at least one complication by the time an individual has been diagnosed. Herman et al. [3] demonstrated a 29% relative risk reduction (RRR) in cardiovascular disease (CVD) outcomes and a 17% RRR in all-cause mortality after 5 years among screened individuals undergoing routine care compared with a 3-year delay in diagnosis and treatment illustrating the beneficial effects accrued from early diagnosis and identification of high-risk individuals. Furthermore, screening for glucose intolerance in high-risk populations and implementing interventions have been shown to be cost-effective [4].

Finally, from 2007 to 2012, approximately 7.9 million individuals in the USA with diabetes were unaware of their diagnosis although 85% had access to a care provider [5]. Therefore, identifying high-risk individuals at an earlier time point is of paramount importance.

Managed by Antonio Secchi.

✉ Michael Bergman
michael.bergman@nyumc.org

¹ Hubert Department of Global Health, Rollins School of Public Health, Emory University, 18, Atlanta, GA, USA

² Department of Endocrinology and Diabetology, Université Catholique de Louvain, University Clinic Saint-Luc, Brussels, Belgium

³ Oporto Medical School, Oporto University, Oporto, Portugal

⁴ NYU Langone Diabetes Prevention Program, Division of Endocrinology and Metabolism, Department of Medicine, NYU School of Medicine, 530 First Avenue, Schwartz East, Suite 5E, New York, NY 10016, USA

Caveats in diagnosing dysglycemic states

The dysglycemic or prediabetic conditions referred to as categories of increased risk for future development of T2DM include impaired fasting glucose (IFG) and IGT.

Defining prediabetes is controversial as different glucose and HbA1c criteria have been posited by the American Diabetes Association (ADA), International Expert Committee (IEC) and World Health Organization (WHO) with differing thresholds, sensitivities, specificities, morbidity and mortality hazard ratios [6] (Table 1). The various definitions therefore identify different but overlapping populations. These inadequacies led to a recent commentary underscoring the need for a more precise, evidence-based definition of intermediate hyperglycemia [6].

Current diagnostic thresholds for T2DM [FPG > 126 mg/dl (7.0 mmol/l) and 2-h plasma glucose > 200 mg/dl (11.1 mmol/l)] are largely based on an association of cross-sectional glycemic levels with diabetic retinopathy [7]. The International Expert Committee (IEC) [8], the ADA [9] and WHO [10] proposed a HbA1c threshold value of 6.5% (47.5 mmol/mol) for diagnosing diabetes also based on studies demonstrating an association between HbA1c and diabetic retinopathy. Discordance between HbA1c and glycemic measures has been observed as they represent different physiologic processes [11]. For example, the US National Health and Nutrition Examination Survey (NHANES-3) from 2005 to 2006 showed that the HbA1c \geq 6.5% (47.5 mmol/mol) identified one-third fewer cases of undiagnosed diabetes than a FPG \geq 126 mg/dl (7.0 mmol/l) [12].

The risk of future outcomes across different prediabetes definitions based on fasting glucose concentration, HbA1c, and 2-h glucose concentration was assessed in the community-based Atherosclerosis Risk in Communities (ARIC) study. The WHO fasting glucose concentration threshold [110–125 mg/dl (6.1–6.9 mmol/l)] and HbA1c-based definitions of prediabetes resulted in lower prevalence estimates than ADA fasting glucose concentration [100–125 mg/dl (5.6–6.9 mmol/l)] and ADA and WHO 2-h glucose concentration cutoffs [140–199 mg/dl

(7.8–11.0 mmol/l)]. However, these were more specific in identifying people at risk for long-term outcomes. ADA fasting glucose and ADA and WHO 2-h glucose definitions of prediabetes were found to be more sensitive for long-term outcomes. Furthermore, HbA1c-based definitions of prediabetes showed stronger associations with long-term outcomes for many major clinical complications. Differences between definitions using both ADA and WHO fasting glucose concentrations for long-term risk associations were not observed when compared with ADA and WHO 2-h glucose concentration cutoffs [13]. The ARIC study also demonstrated that the glycated hemoglobin, in addition to being associated with risk of diabetes, was more strongly associated with risks of cardiovascular disease and death from any cause compared with fasting glucose [13].

Gaps in identifying individuals at high risk for progression to prediabetes and diabetes

Identifying the earliest moment in which to intervene to avert progression to prediabetes and diabetes in high-risk individuals is a substantial challenge. As β -cell function is already substantially impaired in prediabetes based on current definitions [14–17], attention should therefore be focused on identifying individuals with prediabetes even earlier [18]. Biomarkers to monitor progression and identify the time point at which β -cell dysfunction occurs are therefore critically needed [19, 20]. Fasting and post-load glucose levels increased as early as 13 years before diabetes developed in the Whitehall II Study although remaining within the normal range until 2–6 years before diagnosis at which time these levels increased dramatically [21]. Insulin sensitivity was reduced for 13 years and fell steeply 5 years before diagnosis of diabetes. Insulin secretion remained constant until substantial compensation occurred 3–4 years prior to a steep decline prior to diagnosis.

Hence, β -cell function declines in the prediabetic state many years before diabetes develops. As fasting and postprandial glucose concentrations follow a continuum, it is plausible that lifestyle intervention initiated before current

Table 1 Different classifications of prediabetes. Adapted from Ref. [6]

	Impaired fasting glucose	Impaired glucose tolerance	Sub-diabetic HbA1c
ADA (2010)	100–125 mg/dl (5.6–6.9 mmol/L)	140–199 mg/dl (7.8–11.0 mmol/L)	5.7–6.4% (39–46 mmol/mol)
WHO (1999)	110–125 mg/dl (6.1–6.9 mmol/L)	140–199 mg/dl (7.8–11.0 mmol/L)	–
IEC (2009)	–	–	6.0–6.4% (42–46 mmol/mol)

ADA American Diabetes Association, WHO World Health Organization, IEC International Expert Committee

absolute thresholds for prediabetes are achieved and when β -cell function is likely more intact it could be even more effective in preventing progression to diabetes. In fact, lack of progression during the Diabetes Prevention Program (DPP) was related to frequency of regression to normoglycemia [22]. Furthermore, lifestyle intervention was significantly more successful in those with lower baseline glucose concentrations [23]. Use of absolute thresholds for diagnosing subtle dysglycemic states may therefore have the unintended consequence of limiting detection and intervention which may make reversibility less likely [23].

Current absolute definitions of prediabetes remain inadequate as these identify individuals rather late in the dysglycemic continuum, thereby missing an opportunity for earlier intervention when β -cell functionality is more intact [19, 23]. Clearly, harmonization would be contingent on identifying a biomarker characterized by optimal specificity and sensitivity, the ability to diagnose high-risk individuals early in the trajectory to diabetes before β -cell functionality is substantially impaired, as well as the capability of predicting progression to T2DM, complications and mortality. This review will focus on the 1-h post-load plasma glucose during the OGTT as a biomarker fulfilling these characteristics.

The 1-h post-load plasma glucose during the OGTT

Evidence from large-scale population studies has consistently shown that the 1-h plasma glucose (1-h PG) ≥ 155 mg/dl (8.6 mmol/l) during the OGTT may detect incident T2DM and associated complications better than FPG or 2-h PG levels. Published studies investigating the 1-h PG studies are summarized in Table 2. Abdul Ghani et al. [24] demonstrated the predictive power of the 1-h PG level versus FPG and 2-h PG values with incident diabetes over 8 years in a high-risk Mexican–American cohort. They also recommended the 1-h PG threshold of ≥ 155 mg/dl (8.6 mmol/l) and ATP III criteria for the metabolic syndrome in order to stratify high-risk individuals [24]. The Botnia Study and the Malmö Preventive Project provided evidence that fasting glucose, 2-h PG and glucose tolerance status were less efficient predictors than 1-h PG of incident T2DM. The authors concluded that the 1-h PG was a more efficient screening tool to select high-risk individuals for developing T2DM risk [25].

A series of sub-analyses from the CATAnzaro MEtabolic RiSk factors (CATAMERI) study provided further novel insights into the 1-h PG on diabetes and cardiovascular risk factors. The study showed that individuals with NGT and the 1-h PG ≥ 155 mg/dl (8.6 mmol/l) were predisposed to an increased risk for developing diabetes over a 5-year period [26], chronic kidney disease [27], increased risk of

non-alcoholic fatty liver disease diagnosed by ultrasonography [28] and increased vascular stiffness [29] than those with IFG or NGT and 1-h PG < 155 mg/dl (8.6 mmol/l). Mechanistically, sub-analyses also showed an association between NGT individuals and 1-h PG ≥ 155 mg/dl (8.6 mmol/l) with elevated liver enzymes [30], adverse atherogenic profile [31, 32], lower vitamin D concentrations [33], decreased insulin clearance [34], insulin sensitivity and reduced β -cell function [35] and unfavorable inflammatory profile [36] (Table 3).

The accuracy of HbA1c and the 1-h PG versus the OGTT was compared in a real-life clinical setting and found that the level of agreement was twofold greater for the 1-h PG ≥ 155 mg/dl [8.6 mmol/l (95% CI): 0.40(0.28–0.53)] than HbA1c categories defined by the ADA [HbA1c: 5.7–6.4% (39–46 mmol/mol); 0.1(0.03–0.16)] and the IEC [HbA1c: 6.0–6.4% (42–46 mmol/mol); 0.17(0.04–0.30)] [37]. Importantly, the 1-h PG showed a stronger association with 2-h PG, insulin sensitivity index and β -cell function than HbA1c ($P < 0.05$). The Israel Study of Glucose Intolerance, Obesity and Hypertension (The Israel GOH Study) observational study cohort followed over 24 years demonstrated that individuals with a 1-h PG ≥ 155 mg/dl (8.6 mmol/l) but with 2-h PG < 140 mg/dl (7.8 mmol/l) had a significantly elevated risk of both diabetes (OR: 4.35, 95%CI 2.50–7.73) and prediabetes (OR = 1.87, 95%CI 1.09–3.26) after adjusting for sex, age, smoking, body mass index, blood pressure, fasting blood glucose and insulin [38]. In the same cohort, the 1-h PG > 155 mg/dl (8.6 mmol/l) was found to predict mortality even when the 2-h PG was < 140 mg/dl (7.8 mmol/l) after adjusting for sex, age, smoking, body mass index, systolic and diastolic blood pressure. Risk of both progression and mortality were even greater in those with *both* an elevated 1-h PG *and* IGT [39]. The Malmö Preventive Project similarly found increased risk of progression to diabetes, microvascular disease and mortality during a 39-year follow-up in those with a 1-h ≥ 155 mg/dl (8.6 mmol/l) [40]. Therefore, these observations would suggest that the 1-h PG > 155 mg/dl (8.6 mmol/l) may be important for the earlier detection of high-risk individuals to avert T2DM and complications. However, precise estimates regarding how much earlier lifestyle intervention could be initiated with the new 1-h PG definition are difficult to assess in the absence of a prospective trial specifically addressing this question.

Conclusions

As dysglycemic conditions are continuous and cannot be defined by absolute thresholds [23], current diagnostic parameters largely preclude early identification of individuals at high risk for progressing to T2DM. Therefore,

Table 2 Summary of 1-h plasma glucose publications

Author (Year)	Country	Cohort characteristics			Study objectives		Findings
		Sample size (n)	Age (Y)	Male (n, %)	Prevalence 1-h PG > 155 mg/dl (n, %)	BMI (kg/m ²)	
Vaccaro (1992) [42]	USA	873	51.1 ± 7.7	873	NA	27.1 ± 3.5	Examining the relationship 1 h post-load versus 2 h post-load PG measurements to mortality Chicago People Gas Company study; mean follow-up: 19 years Outcome: coronary artery disease 1-h PG _{>195mg/dl} versus 1-h PG _{<100mg/dl} HR [95%CI]: 1.76[0.97–3.20] Outcome: cardiovascular disease 1-h PG _{>195mg/dl} versus 1-h PG _{<100mg/dl} HR [95%CI]: 1.96[1.11–3.45] Outcome: all causes mortality 1-h PG _{>195mg/dl} versus 1-h PG _{<100mg/dl} HR [95%CI]: 1.56[1.06–2.30] Outcome: Diabetes (a) without metabolic syndrome NGT _{1-h PG > 155mg/dl} —metabolic syndrome vs. NGT _{1-h PG < 155 mg/dl} HR [95%CI]: 3.4[1.8–6.4] (b) NGT _{1-h PG > 155 mg/dl} + metabolic syndrome vs. NGT _{1-h PG < 155 mg/dl} HR [95%CI]: 15.2[7.8–29.3]
Abdul Ghani (2008) [24]	USA	1301	42.0 ± 11.0	559 (43)	217 (16.7)	27.1 ± 5.1	Assessing the efficacy of 1-h plasma glucose concentration and the metabolic syndrome in predicting future risk of T2DM San Antonio Heart Study. Mean follow-up duration: 8 years
Cubbedu et al. (2010) [43]	Venezuela	327	39.4 v. 47 (p < 0.05)	Not specified	25 (7.6%)	28.5 v. 29.2 (p < 0.05)	1-h OGTT to predict increased risk of T2DM and surrogate markers of insulin Latino-Hispanics NGT _{1-h PG < 155 mg/dl} vs. NGT _{1-h PG > 155 mg/dl} HOMA-IR: 2.87 v. 2.66 (not significantly different) Matsuda index: 3.42 v. 2.51 (p < 0.05) Disposition index: 0.82 v. 0.49 (p < 0.01)
Manco et al. (2010) [44]	Europe	1100	43.0 v. 45.39 (p < 0.001)	? 56.1% women of total sample would include IGT group	222 (20.2%)	24.9 v. 26.25	To correlate 1-h PG levels with beta cell function RISC study NGT _{1-h PG < 155 mg/dl} vs. NGT _{1-h PG > 155 mg/dl} Beta cell glucose sensitivity: 153.1 v. 86.88 (p < 0.001) B-cell rate sensitivity: 1150.2 v. 781.8 (p < 0.001) Potentiation factor: 2.2 v. 1.9 (p < 0.001)

Table 2 (continued)

Author (Year)	Country	Cohort characteristics			Study objectives		Findings
		Sample size (n)	Age (Y)	Male (n, %)	Prevalence 1-h PG > 155 mg/dl (n, %)	BMI (kg/m ²)	Study design
Marini et al. 2017 [45]	Europe	256	35 v. 40 (<i>p</i> < 0.0001)	100 (39.0%)	72 (28.1%)	28.8 v. 30.1 (<i>p</i> = 0.72)	nondiabetic offspring participating in the EUGENE2 (European Network on Functional Genomics of Type 2 Diabetes) project who had one parent with type 2 diabetes and one parent without a history of type 2 diabetes To study the association between 1-h PG levels and blood viscosity NGT _{1-h} PG < 155 mg/dl vs. NGT _{1-h} PG > 155 mg/dl Matsuda index: 5.89 v. 3.7 (<i>p</i> < 0.0001) Insulin-stimulated glucose disposal: 11.3 v. 8.9 (<i>p</i> = 0.01) Disposition index: 4708 v. 2945 (<i>p</i> < 0.0001)
Succurro et al. (2010) [27]	Italy	1003	49.0 ± 12.0	475 (47.4)	445 (44.3)	30.3 ± 5.9	CATAMERI study cohort. Cross-sectional design Assessing 1-h PG levels as a predictor of CKD Outcome: Chronic kidney disease NGT _{1-h} PG < 155 mg/dl vs. NGT _{1-h} PG > 155 mg/dl OR [95%CI]: 2.61 [1.01–6.77]

Table 2 (continued)

Author (Year)	Country	Cohort characteristics			Study objectives		Findings
		Sample size (n)	Age (Y)	Male (n, %)	Prevalence 1-h PG > 155 mg/dl (n, %)	BMI (kg/m ²)	
Strandberg (2011) [46]	Finland	2756	42	2756	25.7	Helsinki Businessmen Study	Assessing development of diabetes and mortality using the 1-h plasma glucose test in conjunction with BMI
							Outcome: Diabetes. NGT _{1-h} PG>161 mg/dl and BMI ≥ 25 kg/m ² vs. NGT _{1-h} PG<161 mg/dl and BMI < 25 HR [95%CI]: 2.27 [1.67–3.08] NGT _{1-h} PG>161 mg/dl and BMI < 30.0 kg/m ² vs. NGT _{1-h} PG<161 mg/dl and BMI < 25. HR [95%CI]: 4.71 [3.36–6.60] NGT _{1-h} PG > 161 mg/dl and BMI ≥ 30.0 kg/m ² vs. NGT _{1-h} PG<161 mg/dl and BMI < 25. HR [95%CI]: 10.13 [6.46–15.89] Outcome: mortality NGT _{1-h} PG > 161 mg/dl and BMI ≥ 25 kg/m ² vs. NGT _{1-h} PG<161 mg/dl and BMI < 25 HR [95%CI]: 1.14 [1.00–1.29] NGT _{1-h} PG > 161 mg/dl and BMI < 30.0 kg/m ² vs. NGT _{1-h} PG<161 mg/dl and BMI < 25. HR [95%CI]: 1.19 [1.00–1.41] NGT _{1-h} PG>161 mg/dl and BMI ≥ 30.0 kg/m ² vs. NGT _{1-h} PG<161 mg/dl and BMI < 25. HR [95%CI]: 1.51 [1.13–2.02]
Fiorentino (2015) [26]	Italy	234	41.8 ± 10.8	100 (42.9)	60 (25.6)	28.7 ± 5.8	CATAMERI and EUGENE study cohort. Mean follow-up: 5.2 years
							Evaluating 1-h PG levels predicts T2DM in NGT subjects compared to IFG. Outcome: diabetes NGT _{1-h} PG > 155 mg/dl vs. NGT _{1-h} PG < 155 mg/dl HR: 4.02 [1.06–15.26] Outcome: insulin-stimulated glucose disposal: 10.5 v. 8.4 (p < .000 includes isolated IFG and IGT) Disposition index: 265 v. 142 (p < .000 includes isolated IFG and IGT)

Table 2 (continued)

Author (Year)	Country	Cohort characteristics			Study objectives		Findings
		Sample size (n)	Age (Y)	Male (n, %)	Prevalence 1-h PG > 155 mg/dl (n, %)	BMI (kg/m ²)	
Priya (2013) [47]	India	1179	45.4 ± 10.6	734 (62.2)	502 (42.6)	26.3 ± 6.4	<p>Data from primary care; mean follow-up duration: 4.0 years</p> <p>Using the 1-h OGTT to predict progression to prediabetes and T2DM among those with NGT</p> <p>Outcome: Prediabetes NGT_{1-h} PG > 155 mg/dl vs. NGT_{1-h} PG < 155 mg/dl proportion (n, %): 181 (28.9) vs. 211 (52.2)</p> <p>Outcome: diabetes NGT_{1-h} PG > 155 mg/dl vs. NGT_{1-h} PG < 155 mg/dl proportion (n, %): 98 (19.5) vs. 50 (8.0)</p>
Priya et al. (2016) [48]	India	166	20 v. 22 v 23 (p = 0.012)	46.1 v. 66.7 v. 46.7% (p = 0.094)	30 (18.1%)	24.1 v. 25.7 v. 25.2 (p = 0.817)	<p>Data from specialized diabetes center in Chennai</p> <p>Assessing B-cell function and insulin sensitivity indices among NT stratified by 1-h plasma glucose</p> <p>Insulinogenic index: 367 v. 257 v. 246 (p = 0.019)</p> <p>Matsuda index: 11.2 v. 7.4 v. 5.5 (p < 0.001)</p> <p>Insulin sensitivity index: 0.015 v. 0.012 v. 0.011 (p = 0.028)</p>
Kim (2013) [49]	USA	233	11.1 ± 1.7	100 (42.9)	82 (35.2)	28.7 ± 5.5	<p>Obese Latino youth with a family history of type 2 diabetes. Mean follow-up: 4.7 years</p> <p>Examining the utility of elevated 1-h glucose levels to predict deterioration in B-cell function and the development of prediabetes</p> <p>Outcome: prediabetes NGT_{1-h} PG > 155 mg/dl vs. NGT_{1-h} PG < 155 mg/dl OR [95%CI]: 2.4 [1.4–4.2]</p> <p>During the follow-up 1-h PG ≥ 155 mg/dL group exhibited a significantly greater decline in β-cell function compared with youth with a 1-h glucose < 155 mg/dL (β = − 327.8 ± 126.2, p = 0.01)</p>

Table 2 (continued)

Author (Year)	Country	Cohort characteristics			Study objectives		Findings
		Sample size (n)	Age (Y)	Male (n, %)	Prevalence 1-h PG > 155 mg/dl (n, %)	BMI (kg/m ²)	Study design
Oh et al. (2015) [50]	South Korea	192	52.8 v. 53.7	46 (24.0%)	43 (22.4%)	23.8 v. 24.5	Cross-sectional
							To study B-cell function in subjects with 1 h high and 1 h low
							NGT _{1-h} PG < 155 mg/dl vs. NGT _{1-h} PG > 155 mg/dl HOMA-IR: 1.2 v. 1.4 (<i>p</i> = 0.148) Matsuda index: 9.3 v. 8.3 Oral disposition index: 2.5 v. 1.8 (<i>p</i> < 0.001) ISS-2: 301.2 v. 213 (<i>p</i> < 0.001) Insulinogenic index at 60 min was not significantly different between the groups
Bergman et al. (2016) [38, 39]	Israel	739	48.1 ± 6.7	376 (50.9)	188 (25.4)	25.5 ± 3.6	The Israel Study of glucose intolerance, obesity and hypertension. Mean follow-up duration: 24 years
							Evaluating the association of an elevated 1-h PG with cumulative incidence of diabetes, pre-diabetes, and diabetes and mortality
							Outcome: prediabetes 1-h PG _{>155 mg/dl} and 2-h PG _{<140mg/dl} vs. 1-h PG _{<155 mg/dl} and 2-h PG _{<140mg/dl} OR [95%CI]: 1.87 [1.09–3.26] Outcome: diabetes 1-h PG _{>155 mg/dl} and 2-h PG _{<140mg/dl} vs. 1-h PG _{<155 mg/dl} and 2-h PG _{<140mg/dl} OR [95%CI]: 4.35 [2.50–7.73] Outcome: mortality 1-h PG _{>155 mg/dl} and 2-h PG _{<140mg/dl} vs. 1-h PG _{<155 mg/dl} and 2-h PG _{<140mg/dl} OR [95%CI]: 1.28 [1.09–1.50] Outcome: diabetes NGT _{1-h} PG > 145 mg/dl vs. NGT _{1-h} PG < 145 mg/dl HR [95%CI]: 2.84 [2.34–3.45]
Oh et al. (2017) [51]	Korea	5703	51.3 ± 8.7	3062 (53.4)	2435 (42.7)*	24.1 ± 3.1	Korean Genome and Epidemiology Study. Mean follow-up duration: 12 years
							Assessing 1-h OGTT levels as a predictor of future T2DM compared to IGT
Paddock et al. (2017) [52]	USA	1946	25.4	817 (42.0)	NA	30.0 ± 6.0	Southwestern Native American (SWNA) community in Arizona; Mean follow-up duration: 12.8 years
							Assessing 1-h PG as a predictor of T2DM along with insulin secretion
							Outcome: Diabetes NGT _{1-h} PG > 168 mg/dl vs. NGT _{1-h} PG < 168 mg/dl HR [95%CI]: 1.71 [1.60–1.82]

Table 2 (continued)

Author (Year)	Country	Cohort characteristics			Study objectives		Findings
		Sample size (n)	Age (Y)	Male (n, %)	Prevalence 1-h PG > 155 mg/dl (n, %)	BMI (kg/m ²)	
Pareek et al. (2018) [40]	Sweden	4703	48.0 (48.0–49.0)	4703 (100)	1564 (33.3)	24.8 ± 3.1	<p>Evaluating 1-h PG as a predictor of T2DM, cardiovascular risk and mortality compared to 2-h level.</p> <p>Population-based cohort study from Malmö, Sweden. Mean follow-up duration: 39 years</p> <p>Outcome: diabetes NGT_{1-h} PG > 155 mg/dl VS. NGT_{1-h} PG < 155 mg/dl HR [95%CI]: 2.93 [2.48–3.46] Outcome: all-cause mortality NGT_{1-h} PG > 155 mg/dl VS. NGT_{1-h} PG < 155 mg/dl HR [95%CI]: 1.29 [1.19–1.39] Outcome: incident myocardial infarction and fatal ischemic heart disease NGT_{1-h} PG > 155 mg/dl VS. NGT_{1-h} PG < 155 mg/dl HR [95%CI]: 1.24 [1.10–1.39] Outcome: diabetic retinopathy NGT_{1-h} PG > 155 mg/dl VS. NGT_{1-h} PG < 155 mg/dl HR [95%CI]: 5.23 [3.24–8.43] Outcome: diabetic peripheral vascular complications NGT_{1-h} PG > 155 mg/dl VS. NGT_{1-h} PG < 155 mg/dl HR [95%CI]: 2.90 [1.69–4.99]</p>

Table 3 Associations of the elevated 1-h glucose ≥ 155 mg/dL (8.6 mmol/L) with biologic markers and subclinical target organ damage

Impaired β -cell function
Insulin resistance
Adiposity
High lipid ratios
Increased WBC
Increased fibrinogen
Subclinical target organ damage
Subclinical atherosclerosis
Subclinical inflammation
Arterial stiffness
Carotid intima-media thickness
Left ventricular hypertrophy
Decline in kidney function
Fatty liver

prediabetes is diagnosed when β -cell dysfunction is proximal to the development of T2DM. A shift to diagnosing high-risk individuals even earlier offers the potential opportunity for further reducing progression to diabetes, development of microvascular complications and mortality, thereby advancing benefit beyond what has been demonstrated in global diabetes prevention programs. The considerable consistent and significant epidemiologic evidence from different populations substantiates the conclusion that an elevated 1-h PG level appears to be a better alternative for identifying high-risk individuals at a stage when β -cell function is substantially more intact than in prediabetes.

The 1-h PG level has been associated with adverse biologic properties and has been shown to be a marker for subclinical target organ damage (Table 3). Furthermore, as discussed earlier, the 1-h PG possesses desirable characteristics for an optimal biomarker. In addition, shortening the OGTT to 1 h should facilitate its use in clinical practice to avoid underdiagnosing high-risk individuals. Therefore, the aggregate of findings support the proposal that a 1-h PG level ≥ 155 mg/dl (8.6 mmol/l) should be considered for adoption into clinical practice as to earlier detect progression to worsening dysglycemia and mortality [38–40].

Finally, since any single biomarker, including the 1-h PG ≥ 155 mg/dl (8.6 mmol/l), may have limitations, combining biomarkers may provide better sensitivity and specificity for more precisely detecting those at high risk for developing dysglycemia. Additional comparison studies of biomarkers will therefore be required to ascertain their clinical utility [41].

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Human and animal rights This article does not contain any studies with human subjects performed by any of the authors.

Informed consent Informed consent not applicable.

References

1. IDF Atlas-8th edition. <http://www.diabetesatlas.org/key-messages.html>
2. Gregg EW, Zhuo X, Cheng YJ, Albright AL, Venkat Narayan KM, Thompson TJ (2014) Trends in lifetime risk and years of life lost due to diabetes in the USA, 1985–2011: a modeling study. *Lancet Diabetes Endocrinol* 2:867–874
3. Herman WH, Ye W, Griffin SJ et al (2015) Early detection and treatment of type 2 diabetes reduce cardiovascular morbidity and mortality: a simulation of the results of the Anglo-Danish-Dutch study of intensive treatment in people with screen-detected diabetes in primary care (ADDITION-Europe). *Diabetes Care* 38:1449–1455
4. The Diabetes Prevention Program Research Group (2012) The 10-year cost-effectiveness of lifestyle intervention or metformin for diabetes prevention. An intent-to-treat analysis of the DPP/DPPOS. *Diabetes Care* 35:723–730
5. Ali MK, Bullard KM, Gregg EW, del Rio C (2014) A cascade of care for diabetes in the United States: visualizing the gaps. *Ann Int Med* 161(10):681–689
6. Makaroff LE (2017) The need for international consensus on pre-diabetes. *Lancet Diabetes Endocrinol* 5(1):5–7
7. American Diabetes Association (2014) Diagnosis and classification of diabetes mellitus. *Diabetes Care* 37(supplement 1):S81–S90
8. International Expert Committee (2009) International Expert Committee report on the role of the A1C assay in the diagnosis of diabetes. *Diabetes care* 32(7):1327–1334
9. American Diabetes Association (2010) Diagnosis and classification of diabetes mellitus. *Diabetes Care* 33(Supplement 1):S62–S69
10. World Health Organization (2011) Use of glycated haemoglobin (HbA1c) in diagnosis of diabetes mellitus: abbreviated report of a WHO consultation
11. Herman WH, Cohen RM (2012) Racial and ethnic differences in the relationship between HbA1c and blood glucose: implications for the diagnosis of diabetes. *J Clin Endocrinol Metab* 97(4):1067–1072
12. Olson DE, Rhee MK, Herrick K, Ziemer DC, Twombly JG, Phillips LS (2010) Screening for diabetes and pre-diabetes with proposed A1C-based diagnostic criteria. *Diabetes Care* 33(10):2184–2189
13. Warren B, Pankow JS, Punjabi NM et al (2017) Comparative prognostic performance of definitions of prediabetes: a prospective cohort analysis of the Atherosclerosis Risk in Communities (ARIC) study. *Lancet Diabetes Endocrinol* 5:34–42
14. Halban PA, Polonsky KS, Bowden DW et al (2014) β -cell failure in type 2 diabetes: postulated mechanisms and prospects for prevention and treatment. *Diabetes Care* 37(6):1751–1758
15. Zimmet PZ (1992) Kelly West Lecture 1991 challenges in diabetes epidemiology—from West to the rest. *Diabetes Care* 15(2):232–252
16. Lorenzo C, Williams K, Haffner S (2012) Insulin secretion based on the late oral glucose tolerance test period and incident diabetes: the San Antonio Heart Study. *Diabet Med* 29(8):e151–e158
17. Hanson RL, Pratley RE, Bogardus C et al (2000) Evaluation of simple indices of insulin sensitivity and insulin secretion for use in epidemiologic studies. *Am J Epidemiol* 151(2):190–198

18. Jagannathan R, Sevvick MA, Li H et al (2016) Elevated 1-hour plasma glucose levels are associated with dysglycemia, impaired beta-cell function, and insulin sensitivity: a pilot study from a real world health care setting. *Endocrine* 52(1):172–175
19. Bergman M, Dankner R, Roth J, Narayan KV (2013) Are current diagnostic guidelines delaying early detection of dysglycemic states? Time for new approaches. *Endocrine* 44(1):66–69
20. Skyler JS, Bakris GL, Bonifacio E et al (2017) Differentiation of diabetes by pathophysiology, natural history, and prognosis. *Diabetes* 66(2):241–255
21. Tabák AG, Jokela M, Akbaraly TN, Brunner EJ, Kivimäki M, Witte DR (2009) Trajectories of glycaemia, insulin sensitivity, and insulin secretion before diagnosis of type 2 diabetes: an analysis from the Whitehall II study. *Lancet* 373(9682):2215–2221
22. Perreault L, Pan Q, Mather KJ et al (2012) Effect of regression from prediabetes to normal glucose regulation on long-term reduction in diabetes risk: results from the Diabetes Prevention Program Outcomes study. *Lancet* 379(9833):2243–2251
23. Bergman M (2010) Inadequacies of absolute threshold levels for diagnosing prediabetes. *Diabetes/Metab Res Rev* 26(1):3–6
24. Abdul-Ghani MA, Abdul-Ghani T, Ali N, DeFronzo RA (2008) One-hour plasma glucose concentration and the metabolic syndrome identify subjects at high risk for future type 2 diabetes. *Diabetes Care* 31(8):1650–1655
25. Alyass A, Almgren P, Akerlund M et al (2015) Modelling of OGTT curve identifies 1 h plasma glucose level as a strong predictor of incident type 2 diabetes: results from two prospective cohorts. *Diabetologia* 58(1):87–97
26. Fiorentino TV, Marini MA, Andreozzi F et al (2015) One-hour postload hyperglycemia is a stronger predictor of type 2 diabetes than impaired fasting glucose. *J Clin Endocrinol Metab* 100(10):3744–3751
27. Succurro E, Arturi F, Lugara M et al (2010) One-hour postload plasma glucose levels are associated with kidney dysfunction. *Clin J Am Soc Nephrol* 5(11):1922–1927
28. Sesti G, Hribal ML, Fiorentino TV, Sciacqua A, Perticone F (2014) Elevated 1 h postload plasma glucose levels identify adults with normal glucose tolerance but increased risk of non-alcoholic fatty liver disease. *BMJ Open Diabetes Res Care* 2(1):e000016
29. Sciacqua A, Maio R, Miceli S et al (2012) Association between one-hour post-load plasma glucose levels and vascular stiffness in essential hypertension. *PLoS One* 7(9):e44470
30. Succurro E, Arturi F, Grembiale A et al (2011) One-hour post-load plasma glucose levels are associated with elevated liver enzymes. *Nutr Metab Cardiovasc Dis* 21(9):713–718
31. Andreozzi F, Mannino GC, Perticone M, Perticone F, Sesti G (2016) Elevated 1-h post-load plasma glucose levels in subjects with normal glucose tolerance are associated with a pro-atherogenic lipid profile. *Atherosclerosis* 256:15–20
32. Fiorentino TV, Sesti F, Andreozzi F et al (2016) One-hour postload hyperglycemia combined with HbA1c identifies pre-diabetic individuals with a higher cardio-metabolic risk burden. *Atherosclerosis* 253:61–69
33. Sciacqua A, Perticone M, Grillo N et al (2014) Vitamin D and 1-h post-load plasma glucose in hypertensive patients. *Cardiovasc Diabetol* 13:48
34. Marini MA, Frontoni S, Succurro E et al (2013) Decreased insulin clearance in individuals with elevated 1-h post-load plasma glucose levels. *PLoS One* 8(10):e77440
35. Marini MA, Succurro E, Frontoni S et al (2012) Insulin sensitivity, beta-cell function, and incretin effect in individuals with elevated 1-h postload plasma glucose levels. *Diabetes Care* 35(4):868–872
36. Sesti G, Fiorentino TV, Succurro E et al (2014) Elevated 1-h post-load plasma glucose levels in subjects with normal glucose tolerance are associated with unfavorable inflammatory profile. *Acta Diabetol* 51(6):927–932
37. Jagannathan R, Sevvick MA, Fink D, et al (2016) The 1-hour post-load glucose level is more effective than HbA1c for screening dysglycemia. *Acta Diabetol* 53:543–550
38. Bergman M, Chetrit A, Roth J, Jagannathan R, Sevvick M, Dankner R (2016) One-hour post-load plasma glucose level during the OGTT predicts dysglycemia: observations from the 24 year follow-up of the Israel study of glucose intolerance, obesity and hypertension. *Diabetes Res Clin Pract* 120:221–228
39. Bergman M, Chetrit A, Roth J, Dankner R (2016) One-hour post-load plasma glucose level during the OGTT predicts mortality: observations from the Israel study of glucose intolerance, obesity and hypertension. *Diabetic Med.* <https://doi.org/10.1111/dme.13116>
40. Pareek M, Bhatt DL, Nielsen ML et al (2017) Cardiovascular and metabolic risk—enhanced predictive capability of a 1-hour oral glucose tolerance test: a prospective population-based cohort study. *Diabetes Care.* <https://doi.org/10.2337/dc17-1351>
41. Dorcelly B, Katz K, Jagannathan R et al (2017) Novel biomarkers for prediabetes, diabetes, and associated complications. *Diabetes Metab Syndr Obes* 10:345–361
42. Vaccaro O, Ruth KJ, Stamler J (1992) Relationship of postload plasma glucose to mortality with 19-year follow-up. Comparison of one versus two plasma glucose measurements in the Chicago Peoples Gas Company Study. *Diabetes Care* 15(10):1328–1334
43. Cubeddu LX, Hoffmann IS (2010) One-hour postload plasma glucose levels, a predictor of additional risk for diabetes: prevalence, mechanisms, and associated cardiovascular and metabolic risk factors in Hispanics. *Metab Syndr Relat Disord* 8(5):395–402
44. Manco M, Panunzi S, Macfarlane DP et al (2010) One-hour plasma glucose identifies insulin resistance and beta-cell dysfunction in individuals with normal glucose tolerance: cross-sectional data from the Relationship between insulin sensitivity and cardiovascular risk (RISC) study. *Diabetes Care* 33(9):2090–2097
45. Marini MA, Fiorentino TV, Andreozzi F et al (2017) Elevated 1-h post-challenge plasma glucose levels in subjects with normal glucose tolerance or impaired glucose tolerance are associated with whole blood viscosity. *Acta Diabetol* 54(8):775–784
46. Strandberg TE, Pienimäki T, Strandberg AY et al (2011) One-hour glucose, mortality, and risk of diabetes: a 44-year prospective study in men. *Arch Intern Med* 171(10):941–943
47. Priya M, Anjana RM, Chivanga FS, Gokulakrishnan K, Deepa M, Mohan V (2013) 1-hour venous plasma glucose and incident prediabetes and diabetes in Asian Indians. *Diabetes Technol Ther* 15(6):497–502
48. Priya MM, Amutha A, Pramodkumar TA et al (2016) Beta-cell function and insulin sensitivity in normal glucose-tolerant subjects stratified by 1-hour plasma glucose values. *Diabetes Technol Ther* 18(1):29–33
49. Kim JY, Goran MI, Toledo-Corral CM, Weigensberg MJ, Choi M, Shaibi GQ (2013) One-hour glucose during an oral glucose challenge prospectively predicts beta-cell deterioration and pre-diabetes in obese Hispanic youth. *Diabetes Care* 36(6):1681–1686
50. Oh TJ, Min SH, Ahn CH et al (2015) Normal glucose tolerance with a high 1-hour postload plasma glucose level exhibits decreased beta-cell function similar to impaired glucose tolerance. *Diabetes Metab J* 39(2):147–153
51. Oh TJ, Lim S, Kim KM et al (2017) One-hour postload plasma glucose concentration in people with normal glucose homeostasis predicts future diabetes mellitus: a 12-year community-based cohort study. *Clin Endocrinol (Oxf)* 86(4):513–519
52. Paddock E, Hohenadel MG, Piaggi P et al (2017) One-hour and two-hour postload plasma glucose concentrations are comparable predictors of type 2 diabetes mellitus in Southwestern Native Americans. *Diabetologia* 60:1704–1711